



VU Research Portal

A universal carbonate ion effect on stable oxygen isotope ratios in unicellular planktonic calcifying organisms

Ziveri, P.; Thoms, S.; Probert, I.; Geisen, M.; Langer, H.

published in

Biogeosciences
2012

DOI (link to publisher)

[10.5194/bg-9-1025-2012](https://doi.org/10.5194/bg-9-1025-2012)

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Ziveri, P., Thoms, S., Probert, I., Geisen, M., & Langer, H. (2012). A universal carbonate ion effect on stable oxygen isotope ratios in unicellular planktonic calcifying organisms. *Biogeosciences*, 9(3), 1025-1032.
<https://doi.org/10.5194/bg-9-1025-2012>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



A universal carbonate ion effect on stable oxygen isotope ratios in unicellular planktonic calcifying organisms

P. Ziveri^{1,2}, S. Thoms³, I. Probert⁴, M. Geisen³, and G. Langer^{1,3,*}

¹Institute of Environmental Science and Technology (ICTA), Universitat Autònoma de Barcelona (UAB), Bellaterra, 08193, Spain

²Cluster Earth & Climate, Department of Earth Sciences, FALW, VU Universiteit Amsterdam, de Boelelaan 1085 1081HV Amsterdam, The Netherlands

³Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

⁴FR2424, CNRS/UPMC, Station Biologique de Roscoff, 29682 Roscoff, France

*present address: Department of Earth Sciences, Cambridge University Downing St., Cambridge CB2 3EQ, UK

Correspondence to: P. Ziveri (patrizia.ziveri@uab.es)

Received: 23 May 2011 – Published in Biogeosciences Discuss.: 1 August 2011

Revised: 16 February 2012 – Accepted: 16 February 2012 – Published: 15 March 2012

Abstract. The oxygen isotopic composition ($\delta^{18}\text{O}$) of calcium carbonate of planktonic calcifying organisms is a key tool for reconstructing both past seawater temperature and salinity. The calibration of paleoceanographic proxies relies in general on empirical relationships derived from field experiments on extant species. Laboratory experiments have more often than not revealed that variables other than the target parameter influence the proxy signal, which makes proxy calibration a challenging task. Understanding these secondary or “vital” effects is crucial for increasing proxy accuracy. We present data from laboratory experiments showing that oxygen isotope fractionation during calcification in the coccolithophore *Calcidiscus leptoporus* and the calcareous dinoflagellate *Thoracosphaera heimii* is dependent on carbonate chemistry of seawater in addition to its dependence on temperature. A similar result has previously been reported for planktonic foraminifera, supporting the idea that the $[\text{CO}_3^{2-}]$ effect on $\delta^{18}\text{O}$ is universal for unicellular calcifying planktonic organisms. The slopes of the $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ relationships range between $-0.0243\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ (calcareous dinoflagellate *T. heimii*) and the previously published $-0.0022\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ (non-symbiotic planktonic foraminifera *Orbulina universa*), while *C. leptoporus* has a slope of $-0.0048\text{‰} (\mu\text{mol kg}^{-1})^{-1}$. We present a simple conceptual model, based on the contribution of $\delta^{18}\text{O}$ -enriched HCO_3^- to the CO_3^{2-} pool in the calcifying vesicle, which can explain the $[\text{CO}_3^{2-}]$ effect on $\delta^{18}\text{O}$ for the different unicellular calcifiers. This approach provides a new insight into

biological fractionation in calcifying organisms. The large range in $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ slopes should possibly be explored as a means for paleoreconstruction of surface $[\text{CO}_3^{2-}]$, particularly through comparison of the response in ecologically similar planktonic organisms.

1 Introduction

Calcification by marine organisms is a key component of the global carbon cycle. Planktonic calcifying organisms are particularly significant since they are responsible for more than 80 % of global CaCO_3 production and a high proportion of their shells is exported to the seafloor to form carbonate sediments (Milliman, 1993).

The unicellular coccolithophore algae, members of the division Haptophyta, are among the major primary producers and calcifiers worldwide. Coccolithophores are present in virtually all photic zone marine environments (Baumann et al., 2005; Thierstein and Young, 2004) and together with planktic foraminifera they dominate carbonate production in the open ocean (Ziveri et al., 2007; Broecker and Clark, 2008). Another group of planktonic calcifying organisms is a monophyletic lineage of peridiniphycean dinoflagellates that live in the upper water column where light is available for photosynthesis and during their life cycle produce cysts that are characterized by the incorporation of calcite in at least one layer of the cyst wall.

The chemical composition of the calcite produced by these organisms (i.e., foraminiferal tests, the coccoliths of coccolithophores and the cysts of calcareous dinoflagellates) provides a unique source of information with regard to paleoclimatological studies (e.g., Erez and Luz, 1983; Stoll and Ziveri, 2004; Katz et al., 2010).

The oxygen isotope composition of carbonate fossils has been used to reconstruct the most sought-after target in paleoceanography, the sea surface temperature (SST). The $\delta^{18}\text{O}$ -temperature relationship is well characterized in foraminifera (Bemis et al., 1998) and has also been studied in coccolithophores (Ziveri et al., 2003 and references therein) and calcareous dinoflagellate cysts (Zonneveld et al., 2007). In addition to the temperature effect, a dependence of oxygen isotope fractionation on the carbonate chemistry of seawater was demonstrated in four planktic foraminiferal species (Spero et al., 1997; Bijma et al., 1999). A decrease in fractionation with increasing carbonate ion concentration was recorded in tests of these species, but the slopes for this relationship differed by a factor of up to two. The more gentle of the slopes was explained in terms of the isotopic composition of the sum of the carbon species in seawater (Zeebe, 1999), but the steeper slope has remained unexplained to date.

The influence of carbonate chemistry on the oxygen isotope fractionation of coccolithophores and calcareous cyst producing dinoflagellates has never been studied. We therefore conducted controlled laboratory experiments using one of the most important calcite producers among the coccolithophores, *Calcidiscus leptoporus* (Ziveri et al., 2007), as well as the most abundant living calcareous dinoflagellate, *Thoracosphaera heimii*. The carbonate chemistry of seawater was altered to determine its impact on oxygen isotope fractionation during calcification. Based on our experimental data, we propose a simple conceptual model that can explain the different $\delta^{18}\text{O}$ slopes in terms of a different contribution of HCO_3^- conversion to CO_3^{2-} to the establishment of supersaturation with respect to calcite in the calcification vesicle. This model is subsequently applied to explain the data previously obtained from studies on planktic foraminifera (Spero et al., 1997; Bijma et al., 1999).

2 Culture experiments

2.1 Experimental design and sampling

Clonal cultures of *Calcidiscus leptoporus* (strains RCC1154, formerly known as AC360 and AS31 from the Alboran Sea off Spain, and RCC1135 formerly known as AC365 and NS6-1 from the South Atlantic off South Africa) and of *T. heimii* (strain RCC1511 formerly known as JF1 from the Pacific Ocean) from the Roscoff Culture Collection, (<http://www.sb-roscoff.fr/Phyto/RCC/>) were grown in sterile filtered (0.2 μm) seawater enriched with 100 μM nitrate and 6.25 μM phosphate and with trace metal and vitamin sup-

plements according to f/2(-Si) (Guillard and Ryther, 1962). The incident photon flux density was 350 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and a 16/8 h light/dark cycle was applied. Experiments were carried out at 20 °C for *C. leptoporus* and 17 °C for *T. heimii*. Cells were acclimated to experimental conditions for approximately 10 generations and grown in dilute batch cultures in triplicate (Langer et al., 2006). Low cell density at harvest (in general less than 6000 cells per ml) resulted in less than 8 % DIC (dissolved inorganic carbon) consumption (i.e., DIC consumed by the cells at the end of experiment) and a shift in pH of no more than 0.06 units.

The carbonate system of seawater can be manipulated in various ways. Changes in atmospheric $p\text{CO}_2$ can be used to alter $[\text{CO}_2]$, pH and DIC, with TA (total alkalinity) remaining constant. Manipulating $p\text{CO}_2$ requires bubbling with CO_2 gas (constantly throughout the experiment if an open system is employed). Alternatively, the addition of the acid HCl or the base NaOH can be used to adjust $[\text{CO}_2]$, pH, and TA, with DIC remaining constant. Both methods, TA manipulation (*C. leptoporus*) and DIC manipulation (*T. heimii*), were employed in this study, in both cases in closed systems (see below). It should be noted that chemical changes imposed by the two methods are of similar magnitude. The $p\text{CO}_2$ range typically used in experiments is ~ 180 to $\sim 1000 \mu\text{atm}$. This range can be covered by either increasing DIC by approximately 15 % or decreasing TA by approximately 15 % (to increase $[\text{CO}_2]$). The change in either DIC or TA is small compared to the changes in CO_2 concentration at the upper limit (increase by a factor of ~ 6), CO_3^{2-} concentration (decrease by a factor of ~ 4), and H^+ concentration (increase by a factor of ~ 5 , i.e., a pH drop of ~ 0.7 units) (for details on the carbonate system refer to Zeebe and Wolf-Gladrow, 2001). Experiments in which DIC is altered should thus be comparable to experiments in which TA is altered (Schulz et al., 2009; Hoppe et al., 2011).

In order to prevent gas exchange with the atmosphere in our experiments, 2.4 L flasks were filled without headspace and closed with Teflon-lined screw caps. Determination of cell density, however, required regular sampling for cell counts, thereby creating a maximum headspace of 6 ml, and resulting in a negligible shift (3 %) in $\text{CO}_2(\text{aq})$ concentration. Samples for alkalinity measurements were filtered (0.6 μm pore size), poisoned with 1 mL of a HgCl_2 solution (35 g l^{-1}) and stored in 300 mL borosilicate flasks at 0 °C. DIC samples were sterile filtered using a syringe filter (0.2 μm) and stored in 13 mL borosilicate flasks free of air bubbles at 0 °C. Total alkalinity was calculated from linear Gran plots (Gran, 1952) after duplicate potentiometric titration (Bradshaw et al., 1981; Brewer et al., 1986) and DIC was measured photometrically (Stoll et al., 2001) in triplicate. Precision of the total alkalinity measurements was $\sim 3 \mu\text{mol L}^{-1}$ and accuracy $\sim 4 \mu\text{mol L}^{-1}$. For DIC, precision was $\sim 4 \mu\text{mol L}^{-1}$ and accuracy $\sim 5 \mu\text{mol L}^{-1}$. The carbonate system was calculated from temperature, salinity, and the concentrations of DIC, total alkalinity and phosphate, using

the CO₂SYS software (Lewis and Wallace, 1998). Equilibrium constants of Mehrbach et al. (1973) refitted by Dickson and Millero (1987) were chosen.

Cell samples for oxygen and carbon isotope analysis were centrifuged in 50 mL Falcon tubes, centrifuged again in 1 mL tubes in order to remove seawater, dried at 60 °C for 48 h, and stored at room temperature. After the last centrifugation step, care was taken to remove as much of the seawater medium as possible in order to minimize contamination due to residual salts. Seawater samples for isotope analysis were sterile filtered (0.2 µm), poisoned with HgCl₂, and stored in gas tight bottles at room temperature prior to analysis. For determination of cell density, samples were taken daily or every other day, stored at 0 °C and counted within 3 h of sampling using a Sedgwick Rafter counting cell.

2.2 Sample preparation and measurement of isotope ratios

The $\delta^{18}\text{O}$ of water samples from the *C. leptoporus* experiments was measured at the Leibniz Laboratory, Kiel University, on a Finnigan Delta E connected with an equilibration bath (Equi). An aliquot of the sample water was isotopically equilibrated with an external CO₂, this CO₂ was extracted, analyzed by mass spectrometer, and $\delta^{18}\text{O}$ calculated from the isotope composition. $\delta^{18}\text{O}$ calibration of the working reference gas was based on VSMOW, with a fine correction derived from analysis of VSMOW, GISP, and SLAP. This calibration was performed once or twice a year. The stability control in routine operation used internal water standards of different isotopic composition. As with the DIC sample series, a second gas standard was measured at the end of each sample batch. The 1 σ standard deviation of water samples was ± 0.05 for $\delta^{18}\text{O}$. The poisoned filtered medium samples from the *T. heimii* culture experiments were measured at the Vrije Universiteit (Amsterdam) using an on-line gas preparation (Finnigan GasBench II). The 1 σ standard deviation of water samples for $\delta^{18}\text{O}$ was ± 0.05 . Reproducibility tests were based on a routinely run laboratory water standard (Vienna-Standard Mean Ocean Water (V-SMOW)). $\delta^{18}\text{O}$ values ranged from -0.33 to 0.09 ‰ (SMOW).

The stable isotopes of coccolith and calcareous dinoflagellate samples were measured at the Vrije Universiteit (Amsterdam) using a CARBO-KIEL automated carbonate preparation device linked on-line to a Finnigan MAT252 mass spectrometer (Kiel device). The reproducibility of a routinely analyzed carbonate standard (GICS) was better than 0.15 ‰ for $\delta^{18}\text{O}$ for the MAT252. Based on replicate analyses of splits of culture samples, the mean reproducibility of the $\delta^{18}\text{O}$ measurements was better than ± 0.2 ‰.

3 Experimental results and discussion

Carbonate ion concentration in the seawater medium in which the cells were grown, calculated using CO₂SYS (Lewis and Wallace, 1998), ranged from 78 to $530 \mu\text{mol kg}^{-1}$ for *C. leptoporus* experiments and from 117 to $239 \mu\text{mol kg}^{-1}$ for *T. heimii* experiments. The $\delta^{18}\text{O}$ in coccoliths of *C. leptoporus* was inversely correlated with the computed $[\text{CO}_3^{2-}]$ with a regression slope of -0.0048 (‰ $(\mu\text{mol kg}^{-1})$) ($r^2 = 0.89$), (Fig. 1). The $\delta^{18}\text{O}$ in *T. heimii* calcite exhibited the same type of inverse relationship with $[\text{CO}_3^{2-}]$, but with a regression slope of -0.0243 (‰ $(\mu\text{mol kg}^{-1})$) ($r^2 = 0.95$). The 1 σ confidence bounds are shown in Fig. 1. Oxygen isotope fractionation in these two phylogenetically distant unicellular planktonic calcifying taxa, as well as in two planktonic Foraminifera (Spero et al., 1997; Bijma et al., 1999), thus consistently decreases with increasing $[\text{CO}_3^{2-}]$ of seawater. This relationship suggests a common oxygen isotope fractionation mechanism. We propose below a conceptual model that can explain the different slopes identified in this and previous studies based on the contribution of HCO_3^- to the CO_3^{2-} pool in the calcifying vesicle.

The slopes of the $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ relationships for tested planktonic calcifiers range from -0.0022 ‰ $(\mu\text{mol kg}^{-1})^{-1}$ for the foraminifer *Orbulina universa* to -0.0243 ‰ $(\mu\text{mol kg}^{-1})^{-1}$ for the calcareous dinoflagellate *T. heimii* (Fig. 1). *T. heimii* has a steep slope which suggests strong biological control on oxygen isotope fractionation. Of the species analyzed to date, *T. heimii* shows the largest dissimilarity in $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ slope between biological and inorganic precipitates (Spero et al., 1997; Zeebe, 1999). The heavily calcified coccolithophore *C. leptoporus* has a $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ slope of -0.0048 ‰ $(\mu\text{mol kg}^{-1})^{-1}$, similar to that for inorganic precipitates and for the planktonic foraminifer *G. bulloides*, but steeper than the slope reported for *O. universa* (Spero et al., 1997).

In coccolithophores, precipitation of calcite takes place intracellularly within a calcifying vesicle, a membrane-delimited space that is completely isolated from the cytoplasm (e.g., Young et al., 1999). While the pH in the external medium can change under different environmental conditions, cellular pH-homeostasis should keep the pH inside the vesicle constant at an alkaline value favorable for calcite precipitation.

Interestingly, a similar vesicle-based calcification mechanism has been proposed for the common calcareous dinoflagellate *T. heimii* (Inouye and Pienaar, 1983). Although planktic foraminifera are thought to calcify in an extracellular space (Spero, 1988), the role of seawater endocytosis in the biomineralization process in calcareous foraminifera (Bentov et al., 2009) is indeed remarkably similar to the one of coccolithophores and *T. heimii*. This is supported by Mg/Ca




Fig. 1. Effect of $[\text{CO}_3^{2-}]$ on $\delta^{18}\text{O}$ ($\delta^{18}\text{O}_{\text{calcite}} - \delta^{18}\text{O}_{\text{water}}$) values of *C. leptoporus* (coccolithophore), *T. heimii* (calcareous dinoflagellate), *O. universa* (HL = foraminifera grown under photosynthetic maximum light levels, D = maintained in the dark) (Spero et al., 1997), and *G. bulloides* (Spero et al., 1997) (lines are linear regressions fitted to the data). The regression slopes are $-0.0022\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ for *O. universa* (HL = high light; DC = dark conditions), $-0.0048\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ for *C. leptoporus*, $-0.0047\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ for *G. bulloides*, and $-0.024\text{‰} (\mu\text{mol kg}^{-1})^{-1}$ for *T. heimii*. 1σ confidence bounds are shown for each regression (shaded areas).

results in *T. heimii* presented in Gussone et al. (2010) showing strong resemblance with values recorded in planktonic foraminifera thus, supporting the similarities in calcification mechanisms proposed by our model. The extracellular calcification space of foraminifera is isolated from the seawater by a so called pseudopodial network, so that, in effect, also foraminifera calcify in a space which is isolated by means of plasmamembrane not only from the seawater, but also from the cytoplasm (Erez, 2003). This common basic feature of the calcification of the three phylogenetically distinct groups of calcifiers, coccolithophores, foraminifera, and dinoflagellates, can partly account for the fact that it is possible to formulate one single model explaining the dependency of $\delta^{18}\text{O}$ on carbonate chemistry as will be discussed below.

Zuddas and Mucci (1998) and Kim et al. (2006) demonstrated that for an alkaline pH, the kinetics of CaCO_3 precipitation are dominated by the reaction: $\text{Ca}^{2+} + \text{CO}_3^{2-} = \text{CaCO}_3$. Assuming that calcium carbonate is precipitated mainly from carbonate, the $\delta^{18}\text{O}$ composition of the

precipitate simply reflects the isotopic composition of CO_3^{2-} in the vesicle. Since at alkaline pH values CO_3^{2-} is the major carbon source for CaCO_3 it seems reasonable to assume that the uptake of both CO_3^{2-} and HCO_3^- from the surrounding medium contributes to the CO_3^{2-} pool in the vesicle. The uptake of CO_3^{2-} is assumed to yield a pool size proportional to $[\text{CO}_3^{2-}]$ in the external medium (ext): $f \times [\text{CO}_3^{2-}]_{\text{ext}}$ with $f \leq 1$. Inorganic carbon transported into the vesicle (V) as HCO_3^- is converted to CO_3^{2-} , and the $[\text{CO}_3^{2-}]_{\text{V}}$ increases until supersaturation with respect to calcite is obtained in the vesicle and calcium carbonate starts to precipitate at $[\text{CO}_3^{2-}]_{\text{V}} = [\text{CO}_3^{2-}]_{\text{sat}}$. The contribution of bicarbonate conversion to the CO_3^{2-} pool in the vesicle is given by: $[\text{CO}_3^{2-}]_{\text{sat}} - f \times [\text{CO}_3^{2-}]_{\text{ext}}$ (Fig. 2). The H^+ generated during CO_3^{2-} formation from HCO_3^- has to be removed from the vesicle to keep the pH value of the calcifying vesicle constant.

Table 1. Experimental data from this study: CO₂ (μatm), CO₃²⁻ (μmol kg⁻¹), Omega calcite saturation, pH (total scale) values, α and ε fractionation factors (calcite-water) and δ¹⁸O_c-δ¹⁸O_w (calcite-water) (PDB).

species	CO ₂ μatm	CO ₃ ²⁻ μmol kg ⁻¹	Omega calcite	pH total scale	α(x-H ₂ O)	ε(x-H ₂ O)	δ ¹⁸ O _c -δ ¹⁸ O _{H₂O} (PDB)
<i>Calcidiscus leptoporus</i>	953.3	78.5	1.95	7.84	1.0278	27.84	-2.66
<i>Calcidiscus leptoporus</i>	914.6	82.6	2.06	7.86	1.0278	27.76	-2.73
<i>Calcidiscus leptoporus</i>	891.6	84.3	2.1	7.87	1.0278	27.79	-2.71
<i>Calcidiscus leptoporus</i>	636.00	103.4	2.51	8	1.0283	28.29	-2.22
<i>Calcidiscus leptoporus</i>	617.00	105.4	2.56	8	1.0283	28.26	-2.25
<i>Calcidiscus leptoporus</i>	619.00	105.6	2.56	8	1.0284	28.35	-2.16
<i>Calcidiscus leptoporus</i>	377.00	161.7	3.92	8.24	1.0277	27.68	-2.81
<i>Calcidiscus leptoporus</i>	379.00	162.8	3.95	8.24	1.0279	27.87	-2.63
<i>Calcidiscus leptoporus</i>	369.00	166	4.03	8.24	1.0278	27.79	-2.71
<i>Calcidiscus leptoporus</i>	350.3	199.7	4.97	8.26	1.0275	27.45	-3.03
<i>Calcidiscus leptoporus</i>	351.1	200.1	4.98	8.26	1.0276	27.61	-2.89
<i>Calcidiscus leptoporus</i>	332.6	207.9	5.17	8.28	1.0277	27.71	-2.79
<i>Calcidiscus leptoporus</i>	262.00	214.1	5.2	8.4	1.0274	27.38	-3.11
<i>Calcidiscus leptoporus</i>	256.00	219.2	5.32	8.4	1.0274	27.39	-3.10
<i>Calcidiscus leptoporus</i>	237.00	233.1	5.66	8.4	1.0274	27.42	-3.07
<i>Calcidiscus leptoporus</i>	104.2	493.8	12.29	8.72	1.0258	25.78	-4.66
<i>Calcidiscus leptoporus</i>	93	530.2	13.19	8.76	1.0261	26.11	-4.34
<i>Thoracosphaera heimii</i>	205.3	239.4	3.74	8.28	1.0273	27.29	-3.19
<i>Thoracosphaera heimii</i>	208	238.2	3.72	8.01	1.0272	27.19	-3.29
<i>Thoracosphaera heimii</i>	439.9	146.4	2.29	8.01	1.0297	29.46	-0.52
<i>Thoracosphaera heimii</i>	464.1	140	2.19	7.99	1.0296	30.04	-1.09
<i>Thoracosphaera heimii</i>	599.4	117.7	1.84	7.9	1.0303	30.33	-0.24
<i>Thoracosphaera heimii</i>	603.2	116.3	1.82	7.89	1.0298	29.82	-0.74
<i>Thoracosphaera heimii</i>	653.8	110.2	1.72	7.86	1.0301	30.10	-0.47

Chemical equilibrium in the intracellular carbon dioxide system is obtained after several seconds (approximately 15 s at seawater pH = 8.2 and 25 °C, Zeebe et al., 1999), which is negligible compared with the 1.4 h (calculated using data in Langer et al., 2006) required for the formation of one coccolith (of *C. leptoporus*). The time to establish δ¹⁸O equilibrium, on the other hand, is one order of magnitude higher than the time required for coccolith formation (see Table 3.3.9 of Zeebe and Wolf-Gladrow, 2001). It is therefore reasonable to assume that the carbon dioxide system in the calcifying vesicle approaches its chemical equilibrium, whereas the CO₃²⁻ in the vesicle will carry the isotopic fingerprint of the inorganic carbon transported from the cytoplasm into the vesicle. Then, the oxygen isotope fractionation factor between the CO₃²⁻ in the vesicle and water is given by:

$$\alpha_{(V-H_2O)} = \left(\alpha_{(CO_3^{2-}-H_2O)} \times f \times [CO_3^{2-}]_{ext} + \alpha_{(HCO_3^- - H_2O)} \times ([CO_3^{2-}]_{sat} - f \times [CO_3^{2-}]_{ext}) \right) / [CO_3^{2-}]_{sat} \quad (1)$$

Two terms contribute to Eq. (1): the first term represents the equilibrium isotopic composition of CO₃²⁻, whereas the second term is the isotopic composition of the ¹⁸O-enriched HCO₃⁻ flux into the calcifying vesicle. Here, we propose an explanation of the different δ¹⁸O/[CO₃²⁻] slopes for unicellular calcifying organisms in terms of dif-

ferent availability of external carbonate ($f \times [CO_3^{2-}]_{ext}$) for the establishment of supersaturation with respect to calcite in the vesicle ($[CO_3^{2-}]_V = [CO_3^{2-}]_{sat}$). The δ¹⁸O of CO₃²⁻ in the vesicle decreases with increasing external carbonate ion concentration since the contribution of bicarbonate conversion ($[CO_3^{2-}]_{sat} - f \times [CO_3^{2-}]_{ext}$) to the CO₃²⁻ pool in the vesicle decreases as $[CO_3^{2-}]_{ext}$ rises. According to this process, the δ¹⁸O slope is influenced by the calcite saturation product (K_{sp}) in the calcifying vesicle: $K_{sp} = [CO_3^{2-}]_{sat} \times [Ca^{2+}]_{sat}$. The δ¹⁸O/[CO₃²⁻] slope becomes steeper with decreasing K_{sp} , which is a function of temperature and to a lesser extent salinity (Mucci, 1983).

It is still a matter of debate which compartment of the cell takes up Ca²⁺ from the cytoplasm. It has been suggested that the peripheral endoplasmic reticulum (ER) could play this role (Berry et al., 2002). Using the seawater saturation product ($K_{sp} = 10^{-6.415} \text{ mol}^{-2} \text{ kg}^{-2}$ at $S = 32$ and $T = 20^\circ\text{C}$) and for $[Ca^{2+}]_{sat}$ a typical value for $[Ca^{2+}]$ in the ER (500 μmol kg⁻¹; Meldolesi and Pozzan, 1998) yields a $[CO_3^{2-}]_{sat}$ in the vesicle of 769 μmol kg⁻¹. The oxygen isotope fractionation factors (α) between DIC species and water can be calculated using the equations derived by Beck et al. (2005). In freshwater at 20 °C (applicable to *C. leptoporus* experiments), the fractionation factors between HCO₃⁻, CO₃²⁻ and water are $\alpha_{(HCO_3^- - H_2O)} = 1.0325$ and $\alpha_{(CO_3^{2-} - H_2O)} = 1.0254$. At 17 °C (*T. heimii*




Fig. 2. Proposed fractionation mechanism. Calcium carbonate is precipitated mainly from carbonate, therefore the $\delta^{18}\text{O}$ composition of the precipitate simply reflects the isotopic composition of CO_3^{2-} in the vesicle. The uptake of both CO_3^{2-} and HCO_3^- from the surrounding medium contributes to the establishment of the CO_3^{2-} pool in the vesicle. The uptake of CO_3^{2-} is assumed to yield a pool size proportional to $[\text{CO}_3^{2-}]$ in the external medium (ext): $f \times [\text{CO}_3^{2-}]_{\text{ext}}$ with $f \leq 1$. The inorganic carbon transported into the vesicle (V) as HCO_3^- is converted to CO_3^{2-} , and the $[\text{CO}_3^{2-}]_V$ increases until supersaturation with respect to calcite is obtained in the vesicle and calcium carbonate starts to precipitate at $[\text{CO}_3^{2-}]_V = [\text{CO}_3^{2-}]_{\text{sat}}$. The contribution of bicarbonate conversion to the CO_3^{2-} pool in the vesicle is given by: $[\text{CO}_3^{2-}]_{\text{sat}} - f \times [\text{CO}_3^{2-}]_{\text{ext}}$.

experiments), the freshwater fractionation factors are $\alpha_{(\text{HCO}_3^- - \text{H}_2\text{O})} = 1.0332$ and $\alpha_{(\text{CO}_3^{2-} - \text{H}_2\text{O})} = 1.0260$. We applied Eq. (1) using the freshwater α factors at 20°C to simulate the $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ slope for *C. leptoporus*. The measured slope of coccolith $\delta^{18}\text{O}$ vs. $[\text{CO}_3^{2-}]$ coincides with the calculated slope for CO_3^{2-} in the vesicle (-0.0048‰ ($\mu\text{mol kg}^{-1}$) $^{-1}$, see Fig. 1). This slope would require $f = 0.53$ using the proposed model. The model also applies to other calcifiers that possess a vesicle-based calcification mechanism, such as the foraminifera *O. universa* and *G. bulloides*. The shallower slope of *O. universa* (-0.0022‰ ($\mu\text{mol kg}^{-1}$) $^{-1}$) indicates that a considerable amount of CO_3^{2-} in the vesicles of *O. universa* is derived from bicarbonate conversion and requires a f factor value of 0.24.

Note that while $\delta^{18}\text{O}$ slopes can be calculated using Eq. (1), absolute $\delta^{18}\text{O}$ values of CO_3^{2-} in the vesicle, and thereby of the precipitated calcite, cannot be compared. The reason for this is that fractionation factors provided by Beck et al. (2005) refer to freshwater, whereas cells were grown at a salinity of 32. However, the seawater saturation product (at $S = 32$) is not necessarily appropriate for understanding calcite precipitation and ^{18}O fractionation in unicellular calcifying marine organisms. Coccolithophores, for in-

stance, exert strong control over the chemical composition of the calcifying fluid. The ionic strength inside the coccolith vesicle is most likely lower than that of seawater (Anning et al., 1996). In fact, the very steep $\delta^{18}\text{O}$ slope observed in culture experiments with *T. heimii* (-0.024‰ ($\mu\text{mol kg}^{-1}$) $^{-1}$) could be explained in terms of a salinity at the site of calcification (S_V) of 14.64, which is less than half of the salinity in seawater. Based on the saturation product ($K_{\text{sp}} = 10^{-6.803} \text{ mol}^{-2} \text{ kg}^{-2}$ at $S_V = 14.64$ and $T = 17^\circ\text{C}$), a $[\text{Ca}^{2+}]_{\text{sat}}$ of $500 \mu\text{mol kg}^{-1}$ would correspond to a $[\text{CO}_3^{2-}]_{\text{sat}}$ in the vesicle of $315 \mu\text{mol kg}^{-1}$. Using this value for $[\text{CO}_3^{2-}]_{\text{sat}}$ in the vesicle and the freshwater α factors at 17° , the slope of calcite $\delta^{18}\text{O}$ vs. $[\text{CO}_3^{2-}]$ can be fitted using Eq. (1) by assuming a maximum access to the external CO_3^{2-} during calcite precipitation in the vesicle, i.e. $f = 1$ (see Fig. 2).

4 Conclusions

In conclusion, the results presented here show that the negative slope of the $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ relationship that has been recorded in planktic foraminifera is also exhibited by two phylogenetically remote groups of unicellular calcifiers, namely coccolithophores and dinoflagellates. The negative slopes are explained for the first time by a conceptual model that takes physiological mechanisms into consideration. According to the model, the dependence of $\delta^{18}\text{O}$ on carbonate chemistry is mediated through a contribution of HCO_3^- to the CO_3^{2-} pool in the calcifying compartment.

Previous attempts have been made to correct the foraminiferal $\delta^{18}\text{O}$ proxy for variations of paleo- $[\text{CO}_3^{2-}]$ using the experimental slopes during time-windows when seawater carbonate chemistry was notably different. The new results demonstrate similar $\delta^{18}\text{O}/[\text{CO}_3^{2-}]$ slopes (0.0048 – 0.0022 ($\mu\text{mol kg}^{-1}$) $^{-1}$) for planktic foraminifera and coccolithophores. A strong isotopic fractionation is indicated by the steep slope recorded for a calcareous dinoflagellate. There is considerable scope for further exploration of oxygen isotope composition as a tool for paleo- $[\text{CO}_3^{2-}]$ reconstruction by comparison of coccolithophore and dinoflagellate taxa that share (or shared) the same habitat. With the proposed model in mind, further comparative studies on other species of these groups of planktonic calcifiers could provide important mechanistic insights into intra- and inter-group similarities and/or differences in calcification processes. Such studies would also help to assess the degree of bias introduced into $\delta^{18}\text{O}$ -based SST reconstructions by past changes in ocean carbonate chemistry.

Acknowledgements. We thank T. Toyofuku, an anonymous reviewer and the assigned CP editor H. Kitazato for their constructive comments. This research is made possible thanks to the support from the EU FP7 MedSeA project (grant agreement no. 265103), the Spanish Ministry of Science and Innovacion cofunded by the European Fund and FP6 MarinERA (CTM2008-04365-E; CGL2009-10806), The Netherlands Organisation for Scientific Research and the German National Science Foundation DFG grant TH744/2-3. This work was funded in part by The European Research Council (ERC grant 2010-NEWLOG ADG-267931 HE).

Edited by: H. Kitazato

References

- Anning, T., Nimer, N., Merrett, M. J., and Brownlee, C.: Costs and benefits of calcification in coccolithophorids, *J. Mar. Syst.*, 9, 45–56, 1996.
- Baumann, K.-H., Andruleit, H., Böckel, B., Geisen, M., and Kinkel, H.: The significance of extant coccolithophores as indicators of ocean water masses, surface water temperature, and paleoproductivity: a review, *Paläontolog. Zeitschr.*, 79/1, 93–112, 2005.
- Beck, W. C., Grossmann, E. L., and Morse, J. W.: Experimental studies of oxygen isotope fractionation in the carbonic acid system at 15 °, 25 °, and 40 °C, *Geochim. Cosmochim. Acta*, 69, 3493–3503, 2005.
- Bentov, S., Brownlee, C., and Erez, J.: The role of seawater endocytosis in the biomineralization process in calcareous foraminifera, *Proceedings of the National Academy of Sciences*, 51, 21500–21504, 2009.
- Bemis, B. E., Spero, H., Bijma, J., and Lea, D.: Reevaluation of the oxygen isotopic composition of planktonic foraminifera: Experimental results and revised paleotemperature equations, *Paleoceanography*, 13, 150–160, 1998.
- Berry, L., Taylor, A. R., Lucken, U., Ryan, K. P., and Brownlee, C.: Calcification and inorganic carbon acquisition in coccolithophores, *Functional Plant Biology*, 29, 289–299, 2002.
- Bijma, J., Spero, H. J., and Lea, D. W.: Reassessing Foraminiferal Stable Isotope Geochemistry: Impact of the Oceanic Carbonate System (Experimental Results), in: *Use of Proxies in Paleoceanography: Examples from the South Atlantic*, edited by: Fischer, G. a. G. W., Springer Verlag, Berlin Heidelberg, 1999.
- Bradshaw, A. L., Brewer, P. G., Shaffer, D. K., and Williams, R. T.: Measurements of total carbon dioxide and alkalinity by potentiometric titration in the GEOSECS program, *Earth Planetary Sc. Lett.*, 55, 99–115, 1981.
- Brewer, P. G., Bradshaw, A. L., and Williams, R. T.: Measurement of total carbon dioxide and alkalinity in the North Atlantic ocean in 1981, in: *The Changing Carbon Cycle – A Global Analysis*, edited by: Trabalka, J. R. and Reichle, D. E., Springer Verlag, New York, 1986.
- Broecker, W. and Clark, E.: Ratio of coccolith CaCO_3 to foraminifera CaCO_3 in late Holocene deep sea sediments, *Paleoceanography*, 24, PA3205, doi:10.1029/2009PA001731, 2009.
- Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep-Sea Res.*, 34, 1733–1743, 1987.
- Erez, J.: The source of ions for biomineralization in foraminifera and their implications for paleoceanographic proxies, *Reviews in Mineralogy and Geochemistry*, 54, 115–149, 2003.
- Erez, J. and Luz, B.: Experimental paleotemperature equation for planktonic foraminifera, *Geochim. Cosmochim. Acta*, 47, 6, 1025–1031, 1983.
- Gran, G.: Determination of the equivalence point in potentiometric titrations of seawater with hydrochloric acid, *Oceanologica Acta*, 5, 209–218, 1952.
- Guillard, R. R. L. and Ryther, J. H.: Studies of marine planktonic diatoms, I, *Cyclotella nanna* (Hustedt) and *Detonula convervacea* (Cleve), *Canadian Journal of Microbiology*, 8, 229–239, 1962.
- Gussone, N., Zonnefeld, K., and Kuhnert, H.: Minor element and Ca isotope composition of calcareous dinoflagellate cysts of *Thoracosphaera heimii*, *Earth Planetary Sc. Lett.*, 289, 180–188, 2010.
- Hoppe, C., Langer, G., and Rost, B.: *Emiliania huxleyi* shows identical responses to elevated $p\text{CO}_2$ in TA and DIC manipulations, *J. Exp. Mar. Biol. Ecol.*, 406, 54–62, doi:10.1016/j.jembe.2011.06.008, 2011.
- Inouye, I. and Pienaar, R. N.: Observations on the life cycle and microanatomy of *Thoracosphaera heimii* (Dinophyceae) with special reference to its systematic position, *S. Afr. J. Bot.*, 2, 63–75, 1983.
- Katz, M. E., Cramer, B. S., Franzese, A., Hönisch, B., Miller, K. G., Rosenthal, Y., and Wright, J. D.: Traditional and emerging geochemical proxies in foraminifera, *The Journal of Foraminiferal Research*, 40, 165–192, 2010.
- Kim, S.-T., Hillaire-Marcel, C., and Mucci, A.: Mechanisms of equilibrium and kinetic oxygen isotope effects in synthetic aragonite at 25 °C, *Geochim. Cosmochim. Acta*, 70, 5790–5801, 2006.
- Langer, G., Geisen, M., Baumann, K.-H., Kläs, J., Riebesell, U., Thoms, S., and Young, J. R.: Species-specific responses of calcifying algae to changing seawater carbonate chemistry, *Geochem. Geophys. Geosyst.*, 7, Q09006, doi:10.1029/2005GC001227, 2006.
- Lewis, E. and Wallace, D. W. R.: Program Developed for CO_2 System Calculations ORNL/CDIAC-105. Carbon Dioxide Information Analysis Centre, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 1998.
- Mehrbach, C., Culbertson, C. H., Hawley, J. E., and Pytkovicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18, 897–907, 1973.
- Meldolesi, J. and Pozzan, T.: The endoplasmic reticulum Ca^{2+} store: A view from the lumen, *Trends Biochem. Sci.*, 23, 10–14, 1998.
- Milliman, J. D.: Production and accumulation of calcium carbonate in the ocean – budget of a nonsteady state, *Global Biogeochem. Cy.*, 7, 927–957, 1993.
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *Am. J. Sci.*, 283, 780–799, 1983.
- Schulz, K. G., Barcelos e Ramos, J., Zeebe, R. E., and Riebesell, U.: CO_2 perturbation experiments: similarities and differences between dissolved inorganic carbon and total alkalinity manipulations, *Biogeosciences*, 6, 2145–2153, doi:10.5194/bg-6-2145-2009, 2009.
- Spero, H. J.: Ultrastructural examination of chamber morphogenesis and biomineralization in the planktonic foraminifer *Orbulina*

- universa, *Mar. Biol.*, 99, 9–20, 1988.
- Spero, H., Bijma, J., Lea, D. W., and Bemis, B. E.: Effect of seawater carbonate concentration on foraminiferal carbon and oxygen isotopes, *Nature*, 390, 497–500, 1997.
- Stoll, M. H. C., Bakker, K., Nobbe, G. H., and Haese, R. R.: Continuous-flow analysis of dissolved inorganic carbon content in seawater, *Anal. Chem.*, 73, 4111–4116, 2001.
- Stoll, H. M. and Ziveri, P.: Coccolithophore-based geochemical proxies, in: *Coccolithophores: From Molecular Processes to Global Impact*, edited by: Thierstein, H. a. Y., Springer Verlag, Germany, 2004.
- Thierstein, H. R. and Young, J. R.: *Coccolithophores: From Molecular Processes to Global Impact*, Springer, Germany, 2004.
- Young, J. R., Davis, S. A., Bown, P. R., and Mann, S.: Coccolith ultrastructure and biomineralisation, *J. Struct. Biol.*, 126, 195–215, 1999.
- Zeebe, R. E.: An explanation of the effect of seawater carbonate concentration on foraminiferal oxygen isotopes, *Geochim Cosmochimica Acta*, 63, 2001–2007, 1999.
- Zeebe, R. E., Wolf-Gladrow, D. A., and Jansen, H.: On the time required to establish chemical and isotopic equilibrium in the carbon dioxide system in sea water, *Mar. Chem.*, 65, 135–153, 1999.
- Zeebe, R. E. and Wolf-Gladrow, D.: *CO₂ in sea water: equilibrium, kinetics, isotopes*, Elsevier Oceanography Series, 65, Amsterdam, 2001.
- Ziveri, P., Stoll, H., Probert, I., Klaas, C., Geisen, M., Ganssen, G., and Young, J.: Stable isotope “vital effects” in coccolith calcite, *Earth Planet. Sci. Lett.*, 210, 137–149, 2003.
- Ziveri, P., de Bernardi, B., Baumann, K.-H., Stoll, H. M., and Mortyn, P. G.: Sinking of coccolith carbonate and potential contribution to organic carbon ballasting in the deep ocean, *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 659–675, 2007.
- Zonneveld, K. A. F., Mackensen, A., and Baumann, K. H.: Stable oxygen isotopes of *Thoracosphaera heimii* (Dinophyceae) in relationship to temperature; a culture experiment, *Mar. Micropaleont.*, 64, 80–90, 2007.
- Zuddas, P. and Mucci, A.: Kinetics of calcite precipitation from seawater: II. The influence of the ionic strength, *Geochim. Cosmochim. Acta*, 62, 757–766, 1998.